

Revista científica CENTROS
15 de enero de 2018 – Vol. 7 No. 1
ISSN: 2304-604X pp 68-73

Recibido: 10/03/17; Aceptado: 30/5/17

Se autoriza la reproducción total o parcial de este artículo, siempre y cuando se cite la fuente completa y su dirección electrónica.

<https://revistas.up.ac.pa/index.php/centros>

indexada en



<http://www.latindex.unam.mx>



<http://miar.ub.edu/issn/2304-604X>



PRIMER REGISTRO DE *Capucina patula* (WALKER, 1871) (BLATTODEA: BLABERIDAE: ZETOBORINAE) PARA PANAMÁ; UNA NUEVA UBICACIÓN PARA COMPRENDER SU DISTRIBUCIÓN

FIRST RECORD OF *Capucina patula* (WALKER, 1871) (BLATTODEA: BLABERIDAE: ZETOBORINAE) FOR PANAMA; A NEW LOCATION TO COMPRISE YOUR DISTRIBUTION

Stéphane De Greef¹ y Julio Cesar Estrada-Álvarez²

1. Cocobolo Nature Reserve, Mamoni Valley, Chepo District, Panamá.

Email: stephane@stephanedegreef.com

2. Museo Universitario de Historia Natural Dr. Manuel M. Villada UAEMex, Inst. Literario 100, Colonia Centro, Toluca, Estado México C.P. 50000. Entomological Research, Metepec, Estado México.

NOTE RESEARCH

Capucina Saussure, 1893 is a monotypic genus of cockroaches of the subfamily Zetoborinae, its only representative *Capucina patula* (Walker, 1871), described with type locality in Chontales, Nicaragua [= *Zetobora patula* Walker, 1871: 8 (D♂(sic) ♀)]

[=*Capucina cucullata* Saussure, 1893: 67 (♀) junior synonym *Sensu* Princis, 1958: 70, same type material] [=*Capucinus cucullatus*(sic) Saussure & Zehntner, 1893: 102; Lam. VI, Fig. 13 mistake].

Capucina patula is distinctive for the delicate structure, greatly flattened form and very broad tegmina and a brown coloration with a white macula on the anterior edge of the pronotum. Its size is about 30 mm of body length (Hebard, 1921a: 147) and in having the pronotum and tegmina covered with a velvety pilosity (Saussure & Zehntner, 1893) (Fig. 1).




Fig. 1, 2. *Capucina patula* (Walker, 1871) 1) Specimen of Panama; 2) Distribution map. Black point represent previous record, red point is a new record.

Fig. 1, 2. *Capucina patula* (Walker, 1871) 1) Specimen of Panama; 2) Distribution map.

Black point represent previous record, red point is a new record.

Roth in 1970 placing this genus within the Phortioecini tribe (vs McKittrick, 1964 in Panchlorinae, Princis 1960 in Laxtinae) for their affinities and male genitalia (see Roth, 1970). *Capucina patula* has discontinuous records in Nicaragua, Costa Rica and Colombia, without records in Panamanian territory previously (Princis. 1963: 158; Roth, 1970: 224, 233).

With a record for Colombia, Muzo, Boyacá (Apolinar 1937; Hebard, 1921 it was thought that this species was in Panamanian territory, without confirming so far.

On 12 January 2017 the co-author Stéphane De Greef found a single specimen of cockroach on the forest floor during a night walk around the Field Station of Cocobolo Nature Reserve (9.2925°N 79.2056°W) in the Mamoni Valley, Chepo District, Panama Province, Panama. Unable to identify the species, he photographed the specimen the next day on a field studio with backlight white background and posted the resulting image on iNaturalist and various Facebook groups to seek identification by other entomologists. On 10 September 2017, three extremely flat nymphs of cockroaches were found under the peeling bark of a dead tree at another site in the secondary rainforest about one kilometre north of the previous location. The nymphs were recorded on video but were not collected on that occasion, but were regularly seen under the same loose bark over the following months.

Previous records (localities amended): **Nicaragua**, Prov. Depto. Chontales (LT) (Maes, 1992a: 23; Saussure & Zehntner, 1893: 102; Walker, 1871: 8). **Costa Rica**, Prov. Alajuela, Loc. Las Delicias, Santa Clara; Prov. Cartago, Loc. Turrialba, Juan Viñas; Prov. Heredia, La Virgen, Parque Nacional Braulio Carrillo; Prov. San José, Loc. San José; Rio Virilla; Prov. Guanacaste, Loc. Tilarán (Biolley, 1900: 48; Fisk, 1971: 441; Rehn, 1903: 286). **Colombia**, Depto. Boyaca, Mpio. Muzo (Apolinar 1937: 135; Hebard, 1921a: 147).

NOTE: The record of Salazar 2001: 43; Fig 3 is doubtful, because the illustration of the pronoto presented it does not correspond to *Capucina*, is more like *Lanxoblatta frater* Hebard, 1933 from Colombia (ver Hebard, 1933: 23; Lam. II, fig.5), the work

Hebard, 1933 is omitted in Salazar 2001.

New record: 1♀ of Panamá, Prov. Panamá, Distr. Chepo, Valle Mamoni, Reserva Natural Cocobolo (*Cocobolo Nature Reserve*) (9.2925°N 79.2056°W, 220m); 12 /Jan./2017; Stéphane De Greef (Fig. 1 Stéphane De Greef). Stored in Ethanol 90% in the field station of Cocobolo Nature Reserve (Fig. 1)

Biogeography: The monotypic genus *Capucina* Saussure, 1893 have a distribution within the provinces of the Caribbean subregion: East of Central America, West of the Isthmus of Panama (This paper), Amazonia, Choco-Magdalena and Norandina (Vélez et al., 2006) (Fig. 2).

REFERENCE.

Apolinar, M. (1937). Especies nuevas y observaciones diversas sobre dermápteros y ortópteros colombianos. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales correspondiente de la Española*, 1(2): 132-137.

Beccaloni, G. W. (2014). Cockroach Species File Online. Version 5.0/5.0. World Wide Web electronic publication. <<http://Cockroach.SpeciesFile.org>> [accessed 11 July 2014].

Biolley, P. (1900). IV Ortopteres recogidos en Costa Rica desde 1890 a 1900. *Informe Museo Nacional de Costa Rica 1899-1900*: 40-49.

Fisk, F.W. (1971). An annotated checklist of Costa Rican cockroaches (Dictyoptera: Blattaria). *Proceedings of the Entomological Society of Washington*, 73(4):431-444.

Hebard, M. (1921). Studies in the Dermaptera and Orthoptera of Colombia. Second Paper. Dermaptera and Orthopterous Families Blattidae, Mantidae and

- Phasmidae. *Transactions of the American Entomological Society*, 47(2): 107-169.
- Hebard, M. (1933). Studies in the Dermaptera and Orthoptera of Colombia. Supplement to Papers One to Five. *Transactions of the American Entomological Society*, 59(1): 13-67.
- Maes, J. M. (1992). Catálogo de los Blattodea (Dictyoptera) de Nicaragua. I. Familias Polyphagidae y Blaberidae. *Revista Nicaragüense de Entomología*, 19: 21-28.
- McKittrick, F. A. (1964). Evolutionary studies of cockroaches. Cornell University Agricultural Experiment Station.
- Princis, K. (1958). Revision der Walkerschen und Kirbyschen Blatterientypen im British Museum of Natural History, London II. *Opuscula Entomologica*. 23:59-75.
- Princis, K. (1960). Zur systematik der Blattarien. *Eos: Revista Española de Entomología* 36(4): 427-449.
- Princis, K. (1963). Blattariae: Suborde [sic] Polyphagoidea: Fam.: Homoeogamiidae, Euthyrrhaphidae, Latindiidae, Anacompsidae, Atticolidae, Attaphilidae. Subordo Blaberoidea: Fam. Blaberidae. In: Beier, M. (Ed.). *Orthopterorum Catalogus*. Pars 4. W. Junk's-Gravenhage: 76-172.
- Rehn, J. A. G. (1903). Studies in American Blattidae. *Transactions of the American Entomological Society*, 29: 259-290.
- Roth, L. M. (1970). The Male Genitalia of Blattaria. III. Blaberidae: Zetoborinae. *Psyche*, 77: 217-236.
- Saussure, H. & L. Zehntner. (1893). Insecta-Orthoptera. Vol. I. *Biologia Centrali-Americana*, Porter (ed.):1-112.

Saussure, H. (1893). De quelques genres de Blattes, tribu des blattiens (parte 2).

Societas Entomologica, 9: 67-68.

Salazar, J. (2001). Blattodea de Colombia. Nuevas adiciones y rectificaciones a los

Mántidos de la primera parte (Insecta: Mantodea). *Boletín Científico Centro de Museos. Museo de Historia Natural*, 5: 38-63.

Vélez, A., M. Wolff & E. Gutiérrez. (2006). Blattaria of Colombia: List and distribution of genera. *Zootaxa* 1210: 39–52

Walker, F. (1871). *Catalogue of the specimens of Dermaptera Saltatoria and supplement of the Blattari in the collection of the British Museum*. Trustees of the British Museum.